

## **Environmental Consequences of Benzo[a]pyrene in Fish**

<sup>1</sup>Shikha Dwivedi

<sup>2</sup>Manoj Kumar

<sup>3</sup>Sunil P. Trivedi

### **Abstract**

Polycyclic aromatic hydrocarbons (PAHs) are widespread contaminants in aquatic environment. Benzo[a]pyrene (B[a]P) is one of the prototypes of PAHs which is formed during the incomplete burning of organic materials. High-performance liquid chromatography (HPLC) and Gas chromatography mass spectrometry (GC-MS) are the appropriate tool to assess B[a]P concentration in fish. This study reveals that the B[a]P accumulation in fish which alters ROS formation, antioxidants activity, MN induction, histological lesions, and molecular mechanisms.

---

**Keywords:** Polycyclic aromatic hydrocarbons, Benzo[a]pyrene, Oxidative stress, Genotoxicity, Fish

---

### **Introduction**

Polycyclic aromatic hydrocarbons (PAHs) are universal pollutants in aquatic habitats. The major portion of PAHs arrives into water bodies from land-based runoff or atmospheric deposition. Though PAH levels deteriorated in urban watersheds from 1970s to 1980s because of decrement in coal burning and industrial discharges (Guntupalli et al., 2016; Parmar et al., 2020; Yuan et al., 2021). But the last time period has formed new increases in PAH accumulation in aquatic organisms due to automobile practice linked through urban extension (Saha et al., 2009) or diesel burning by heavyweight vehicles (Baum et al., 2016). Furthermore,

---

<sup>1</sup> Research Scholar, Environmental Toxicology & Bioremediation Laboratory, Department of Zoology, University of Lucknow, E-mail: [shikha.dwivedi0000@gmail.com](mailto:shikha.dwivedi0000@gmail.com)

<sup>2</sup> Assistant Professor, Environmental Toxicology & Bioremediation Laboratory, Department of Zoology, University of Lucknow

<sup>3</sup>Professor, Environmental Toxicology & Bioremediation Laboratory, Department of Zoology, University of Lucknow, E-mail: [sat060523@gmail.com](mailto:sat060523@gmail.com)

new studies address that petroleum hydrocarbons produced from oil spills can easily persist in aquatic sediments, plants and animals for years or longer (Vanzella et al., 2007).




Figure 1: B[a]P as a toxic substance.

B[a]P is one of the carcinogenic petroleum chemicals with two or more fused aromatic rings which is widely dispersed in the environment (Harris et al., 2020). The production or utilization of B[a]P is not done commercially, but is generally available in the environment. It is formed primarily as outcome of pyrolytic events, particularly the incomplete incineration of organic resources in industrial and various human actions like processing of crude oil and coal, ignition of natural gas for culinary and wood burning (Hamilton et al., 2021; IARC, 2014; Le Bihanic et al., 2014). Additional human actions that result in B[a]P release include the garbage burning, road traffic by vehicles, tobacco and cigarette smoking, and ingestion of charcoal roasted and smoked foods (Ramesh and Archibong, 2011; Sen and Field, 2013). B[a]P binds to small elements in the air and is easily diffused into the environment through the air. B[a]P can then be inhaled or deposited in the environment on plants, soil, and water bodies (Guo et al., 2021; Jabeen et al., 1234; WHO, 2003). Aquatic animals and plants may then come in contact with B[a]P. B[a]P, and its metabolites can persist in the aquatic environment for many years. After the exposure of B[a]P in the residues of ponds, lakes, and waterways, it accumulates in the organs of aquatic invertebrates and vertebrates (Batel et al., 2018; Fanali et al., 2018; Huang et al., 2014). Like other PAHs, B[a]P is mainly toxic to aquatic organism in the occurrence of UV light, increasing health risk to the animals in unshaded or shallow environments (Sen and Field, 2013).

Fishes eagerly take up lipophilic organic substances such as B[a]P from the aquatic environment, with various physiological, cellular and molecular effects (Albornoz-Abud et al., 2021; Honda and Suzuki, 2020). Fish are able to break down xenobiotics, so there is little

accumulation of B[a]P in fish (Scott and Jones, 2000). However, B[a]P acts as a genotoxicant in aquatic organisms, causing many biochemical changes most studied in fish, and the studies on effects of B[a]P on physiological perturbations of fish are comparatively limited. Furthermore, studies are needed to explore the toxic effects of B[a]P in fish.

The quantification of physiological, biochemical, and molecular constraints is a diagnostic implement regularly used in aquatic toxicology and biomonitoring (Dong et al., 2019; Javed and Usmani, 2019). Micronucleus (MN) test is a very reliable tool to assess the genotoxic potential of any xenobiotics (Awasthi et al., 2019). MN formation in erythrocytes of fish by single and double strands break of DNA, and may also a result of the inappropriate DNA repair or failed DNA repair mechanism (Fenech, 2002). Assessment of oxidative stress reveals the redox state of the cell. To deal with enhanced Reactive oxygen species (ROS) in cell, defence mechanism works in which enzymatic- superoxide dismutase (SOD), catalase (CAT), Glutathione reductase (GR), and non-enzymatic antioxidant- reduced glutathione (GSH) assist to maintain the homeostasis in cell (Jifa et al., 2006). The oxidative stress generated DNA damage and alter different molecular mechanisms such as apoptosis, and autophagy in fish. The present study comprises of different techniques for assessing B[a]P toxicity and its deleterious effects in fish.

## **Materials and methods**

### **Estimation of B[a]P in fish**

The B[a]P estimation in fish samples can be done by using High-performance liquid chromatography (HPLC) and Gas chromatography mass spectrometry GC-MS techniques:

#### **Estimation through HPLC**

In order to quantify B[a]P accumulation, fish samples can be obtained using the Soxhlet extraction method with a mixture of acetone and dichloromethane (1:1). The obtained extracts will be dried by anhydrous sodium sulfate and then concentrated through a rotary evaporator. The concentrated extract will then be diluted with 10 mL of hexane and then further concentrated with 2 mL of hexane. The prepared extracts will be used for the analyses of B[a]P in fish tissues using water–acetonitrile solvent organization on HPLC equipped with pump and UV–vis detector. C-18 column will be preferred for the B[a]P detection. The flow rate can be adjusted to 1 mL/min. The B[a]P can be recognized by retention time compared with reference to the B[a]P standards (Malik et al., 2008).

## Estimation through GC-MS

The lyophilized fish samples will be used for B[a]P determination through GC-MS. For GC-MS analysis, carrier gas-Helium and the column pressure-10 psi will be maintained to provide an estimated flow rate of 1 ml/min. The injector lines at 290 °C and transfer lines at 250 °C will be maintained. The temperature of columns will be maintained at °C for 4 min, ramped at 300 °C at a range of 10 °C/min. The mass spectrometer will be employed in electron ionization manner and all the spectra will be obtained with a huge range of m/z 50–400 and automatic gain control (AGC) (Giri et al., 2013).

## Experimental design on fish

Freshwater live of test model will be collected and brought to the laboratory in an aerated medium (Trivedi et al., 2021). During the acclimation, fish will be kept in ventilated glass aquaria and fed with food pellets twice a day. All the necessary physiochemical properties (pH, Temperature, DO, Alkalinity, and Hardness) of aquaria water will be maintained. After the 15 days acclimatization, healthy specimens will be randomly distributed in all the experimental aquaria. The experimental groups were selected in triplicates. Sampling will be done at estimated time interval. Test parameters- Micronucleus test, ROS, biochemical activity (SOD, CAT, GSH and GR), histopathology and transcriptional analysis can be performed at each sampling duration by following the protocol of Trivedi et al., (2022).




Figure 2: Effects of B[a]P.

## B[a]P estimation and its detrimental effects in fish

The measurement of B[a]P and other PAHs in surface water, sediments and fish was performed by using HPLC from eight different locations of the river Gomti (India) (Malik et.

al., 2010 and Malik et al., 2008). Similarly, Palanikumar et al., 2012 investigated the accumulation of B[a]P through GC-MS after its sub-lethal exposure in fish, *Chanos chanos*. B[a]P when enters into the fish, different metabolites formed which cause ROS production via redox cycle. Qi and Tang, (2020) also examined the ROS production after the 96h exposure of B[a]P in mussel *Mytilus coruscus*. Increased ROS production may cause extreme oxidative stress, to deal with this oxidative stress an antioxidant defense system works with different enzymatic (SOD, CAT and GR) and non-enzymatic (GSH) antioxidants. Antioxidant-SOD acts as first line defence against oxidative stress caused due to B[a]P and catalyzes excess superoxide radicals into H<sub>2</sub>O<sub>2</sub> (Gravato and Guilhermino, 2009; Guo et al., 2021). SOD induction was also found in different fish species exposed to waterborne contaminant B[a]P (Dellali et al., 2021; Qi and Tang, 2020). Further, CAT enzyme breaks H<sub>2</sub>O<sub>2</sub> into water and augmented CAT activity designates the higher concentration of H<sub>2</sub>O<sub>2</sub> in cell. Rodrigues et al., (2022) also measured the increased activity of CAT enzyme in tissues of *Scrobicularia plana* after B[a]P exposure. GR maintains the redox potential of cell by regenerating GSH from GSSG. Increased level of GR was documented in tissues of mussel *Mytilus coruscus* exposed to B[a]P (Qi and Tang, 2020). A non-enzymatic antioxidant-GSH was also estimated in different body parts of fish (Jifa et al., 2006; Santos and Bueno, 2020). Studies clarified that the B[a]P was found genotoxic for the fish. It upsurges the frequency of MN formation in cells. After the acute exposure of B[a]P, raised MN frequency were detected in rainbow trout and common carp (Kim and Hyun, 2006). Another study established that B[a]P raised the MN formation in milkfish-*Chanos chanos* after 96 h of experimental duration (Palanikumar et al., 2012).

Histopathological results may help to examine a range of anomalies in the liver and kidney tissues of fish. Numerous studies have been reported the different histological modifications tissues of fish after B[a]P exposure (Briaudeau et al., 2021; Carlson et al., 2004; Esmaeilbeigi et al., 2021; Woo, 2022).

Transcriptional analysis helps to quantify the expression of specific genes. Various studies have been done on mRNA expression in different organs exposed to B[a]P. In a study, up-regulation of tumor suppressor gene-p53 was found involved in autophagy by down-regulating the mTOR gene after B[a]P exposure which is investigated by Lin et al., (2016) and Sforzini et al., (2017).

Indeed, more studies need to be performed to evaluate the toxic ability of B[a]P in different tissues of fish based on MN, antioxidant enzymes, transcriptional analyses, and other related biomarkers.

## Conclusion

The present study reveals the toxic potential of B[a]P in fish and its deleterious effects on genotoxicity, oxidative stress, histopathological impressions and transcriptional analysis in fish. The study also discloses the assessment of B[a]P accumulation and its genotoxic ability in fish. B[a]P is also able to accumulate in fish and cause oxidative stress by producing ROS. Thus, the study helps in understanding the toxic impact of B[a]P in fish and performing the future studies on aquatic creatures. Increasing concentration of B[a]P causing major health risk to aquatic organisms.

## References

- Albornoz-Abud, N.A., Canul-Marín, G.F., Chan-Cuá, I., Hernández-Núñez, E., Cañizares-Martínez, M.A., Valdés-Lozano, D., Rodríguez-Canul, R., Albores-Medina, A., Colli-Dula, R.C., 2021. Gene expression analysis on growth, development and toxicity pathways of male Nile tilapia (*Oreochromis niloticus*), after acute and sub-chronic benzo ( $\alpha$ ) pyrene exposures. *Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology* 250, 1–13. <https://doi.org/10.1016/j.cbpc.2021.109160>
- Awasthi, Y., Ratn, A., Prasad, R., Kumar, M., Trivedi, A., Shukla, J.P., Trivedi, S.P., 2019. A protective study of curcumin associated with Cr6+ induced oxidative stress, genetic damage, transcription of genes related to apoptosis and histopathology of fish, *Channa punctatus* (Bloch, 1793). *Environmental Toxicology and Pharmacology* 71, 1–10. <https://doi.org/10.1016/j.etap.2019.103209>
- Batel, A., Borchert, F., Reinwald, H., Erdinger, L., Braunbeck, T., 2018. Microplastic accumulation patterns and transfer of benzo[a]pyrene to adult zebrafish (*Danio rerio*) gills and zebrafish embryos. *Environmental Pollution* 235, 918–930. <https://doi.org/10.1016/j.envpol.2018.01.028>
- Baum, G., Kegler, P., Scholz-Böttcher, B.M., Alfiansah, Y.R., Abrar, M., Kunzmann, A., 2016. Metabolic performance of the coral reef fish *Siganus guttatus* exposed to combinations of

- water borne diesel, an anionic surfactant and elevated temperature in Indonesia. *Marine Pollution Bulletin* 110, 735–746. <https://doi.org/10.1016/j.marpolbul.2016.02.078>
- Briaudeau, T., Alves Dos Santos, L.A., Zorita, I., Izagirre, U., Marigómez, I., 2021. Biological responses and toxicopathic effects elicited in *Solea senegalensis* juveniles by waterborne exposure to benzo[a]pyrene. *Marine Environmental Research* 170. <https://doi.org/10.1016/j.marenvres.2021.105351>
- Carlson, E.A., Li, Y., Zelikoff, J.T., 2004. Benzo [ a ] pyrene-induced immunotoxicity in Japanese medaka ( *Oryzias latipes* ): relationship between lymphoid CYP1A activity and humoral immune suppression 201, 40–52. <https://doi.org/10.1016/j.taap.2004.04.018>
- Dellali, M., Hedfi, A., Ben, M., Noureldeen, A., Darwish, H., Beyrem, H., Gyedu-ababio, T., Dervishi, A., Karachle, P.K., Boufahja, F., 2021. Comparative Biochemistry and Physiology , Part C decussatus as a predictor of pelago-benthic responses after exposure to Benzo [ a ] Pyrene. *Comparative Biochemistry and Physiology, Part C* 249, 109141. <https://doi.org/10.1016/j.cbpc.2021.109141>
- Dong, J., Wang, J. qin, Qian, Q., Li, G. chun, Yang, D. qin, Jiang, C., 2019. Micronucleus assay for monitoring the genotoxic effects of arsenic in human populations: A systematic review of the literature and meta-analysis. *Mutation Research - Reviews in Mutation Research*. <https://doi.org/10.1016/j.mrrev.2019.02.002>
- Esmailbeigi, M., Kalbassi, M.R., Seyedi, J., Tayemeh, M.B., Moghaddam, J.A., 2021. Intra and extracellular effects of benzo [α] pyrene on liver, gill and blood of Caspian White fish (*Rutilus frissi kutum*): Cyto-genotoxicity and histopathology approach. *Marine Pollution Bulletin* 163, 111942. <https://doi.org/10.1016/j.marpolbul.2020.111942>
- Fanali, L.Z., Franco-Belussi, L., Bonini-Domingos, C.R., de Oliveira, C., 2018. Effects of benzo[a]pyrene on the blood and liver of *Physalaemus cuvieri* and *Leptodactylus fuscus* (Anura: Leptodactylidae). *Environmental Pollution* 237, 93–102. <https://doi.org/10.1016/j.envpol.2018.02.030>
- Fenech, M., 2002. Chromosomal biomarkers of genomic instability relevant to cancer. *Drug Discovery Today* 7, 1128–1137. [https://doi.org/10.1016/S1359-6446\(02\)02502-3](https://doi.org/10.1016/S1359-6446(02)02502-3)
- Fenech, M., 1993. The Cytokinesis-Block Micronucleus Technique and Its Application to Genotoxicity Studies in Human Populations, *Environmental Health Perspectives*

## Supplements.

- Giri, B., Patel, K.S., Jaiswal, N.K., Sharma, S., Ambade, B., Wang, W., Simonich, S.L.M., Simoneit, B.R.T., 2013. Composition and sources of organic tracers in aerosol particles of industrial central India. *Atmospheric Research* 120–121, 312–324. <https://doi.org/10.1016/j.atmosres.2012.09.016>
- González, A., Espinoza, D., Vidal, C., Moenne, A., 2020. Benzopyrene induces oxidative stress and increases expression and activities of antioxidant enzymes, and CYP450 and GST metabolizing enzymes in *Ulva lactuca* (Chlorophyta). *Planta* 252, 1–13. <https://doi.org/10.1007/s00425-020-03508-w>
- Gravato, C., Guilhermino, L., 2009. Effects of benzo(a)pyrene on seabass (*Dicentrarchus labrax* L.): Biomarkers, growth and behavior. *Human and Ecological Risk Assessment* 15, 121–137. <https://doi.org/10.1080/10807030802615659>
- Guntupalli, S., Thunuguntla, V.B.S.C., Santha Reddy, K., Issac Newton, M., Rao, C. V., Bondili, J.S., 2016. Enhanced Degradation of Carcinogenic PAHs Benzo (a) Pyrene and Benzo (k) Fluoranthene by a Microbial Consortia. *Indian Journal of Science and Technology* 9. <https://doi.org/10.17485/ijst/2016/v9i35/93590>
- Guo, B., Feng, D., Xu, Z., Qi, P., Yan, X., 2021. Acute benzo[a]pyrene exposure induced oxidative stress, neurotoxicity and epigenetic change in blood clam *Tegillarca granosa*. *Scientific Reports*. <https://doi.org/10.1038/s41598-021-98354-5>
- Hamilton, T.J., Krook, J., Szaszkievicz, J., Burggren, W., 2021. Shoaling, boldness, anxiety-like behavior and locomotion in zebrafish (*Danio rerio*) are altered by acute benzo[a]pyrene exposure. *Science of the Total Environment* 774, 145702. <https://doi.org/10.1016/j.scitotenv.2021.145702>
- Harris, J.B., Hartman, J.H., Luz, A.L., Wilson, J.Y., Dinyari, A., Meyer, J.N., 2020. Zebrafish CYP1A expression in transgenic *Caenorhabditis elegans* protects from exposures to benzo[a]pyrene and a complex polycyclic aromatic hydrocarbon mixture. *Toxicology* 440, 152473. <https://doi.org/10.1016/j.tox.2020.152473>
- Honda, M., Suzuki, N., 2020. Toxicities of polycyclic aromatic hydrocarbons for aquatic animals. *International Journal of Environmental Research and Public Health* 17. <https://doi.org/10.3390/ijerph17041363>



- Huang, L., Chernyak, S.M., Batterman, S.A., 2014. PAHs (Polycyclic Aromatic Hydrocarbons), Nitro-PAHs, Hopanes and Steranes Biomarkers in Sediments of Southern Lake Michigan, USA. *The Science of the total environment* 487, 173. <https://doi.org/10.1016/J.SCITOTENV.2014.03.131>
- International Agency for Research on Cancer (IARC), 2014. Chemical Agents and Related Occupations. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans 100F, 423–428.
- Jabeen, G., Farkhanda Manzoor, ·, Javid, A., Azmat, H., Arshad, · Mateen, Fatima, S., 1234. Evaluation of Fish Health Status and Histopathology in Gills and Liver Due to Metal Contaminated Sediments Exposure. *Bulletin of Environmental Contamination and Toxicology* 100, 492–501. <https://doi.org/10.1007/s00128-018-2295-7>
- Javed, M., Usmani, N., 2019. An Overview of the Adverse Effects of Heavy Metal Contamination on Fish Health. *Proceedings of the National Academy of Sciences India Section B - Biological Sciences* 89, 389–403. <https://doi.org/10.1007/s40011-017-0875-7>
- Jifa, W., Yu, Z., Xiuxian, S., You, W., 2006. Response of integrated biomarkers of fish (*Lateolabrax japonicus*) exposed to benzo[a]pyrene and sodium dodecylbenzene sulfonate. *Ecotoxicology and Environmental Safety* 65, 230–236. <https://doi.org/10.1016/j.ecoenv.2005.08.002>
- Kim, I.Y., Hyun, C.K., 2006. Comparative evaluation of the alkaline comet assay with the micronucleus test for genotoxicity monitoring using aquatic organisms. *Ecotoxicology and Environmental Safety* 64, 288–297. <https://doi.org/10.1016/j.ecoenv.2005.05.019>
- Le Bihanic, F., Clérandeau, C., Le Menach, K., Morin, B., Budzinski, H., Cousin, X., Cachot, J., 2014. Developmental toxicity of PAH mixtures in fish early life stages. Part II: adverse effects in Japanese medaka. *Environmental Science and Pollution Research* 21, 13732–13743. <https://doi.org/10.1007/s11356-014-2676-3>
- Lin, C., Chen, T., Tseng, Y., Wu, G., Hsieh, M., 2016. Honokiol induces autophagic cell death in malignant glioma through reactive oxygen species-mediated regulation of the p53 / PI3K / Akt / mTOR signaling pathway. *Toxicology and Applied Pharmacology* 304, 59–69. <https://doi.org/10.1016/j.taap.2016.05.018>
- Malik, A., Ojha, A.P., Singh, A.K.P., 2008. Distribution of Polycyclic Aromatic

- Hydrocarbons in Edible Fish from Gomti River , India 134–138.  
<https://doi.org/10.1007/s00128-007-9331-3>
- Palanikumar, L., Kumaraguru, A.K., Ramakritinan, C.M., Anand, M., 2012. Biochemical response of anthracene and benzo [a] pyrene in milkfish *Chanos chanos*. *Ecotoxicology and Environmental Safety* 75, 187–197. <https://doi.org/10.1016/j.ecoenv.2011.08.028>
- Parmar, H., Yusufzai, S., Parmar, P., Bajaniya, V., 2020. Efficacy of Shatavari supplemented diet on growth performance of genetically improved farmed tilapia ( GIFT ) fry. *Journal of Entomology and Zoology Studies* 8, 559–561.
- Qi, P., Tang, Z., 2020. The Nrf2 molecule trigger antioxidant defense against acute benzo ( a ) pyrene exposure in the thick shell mussel *Mytilus coruscus*. *Aquatic Toxicology* 226, 105554. <https://doi.org/10.1016/j.aquatox.2020.105554>
- Ramesh, A., Archibong, A.E., 2011. Reproductive Toxicity of Polycyclic Aromatic Hydrocarbons: Occupational Relevance, Reproductive and Developmental Toxicology. Elsevier Inc. <https://doi.org/10.1016/B978-0-12-382032-7.10043-8>
- Rodrigues, A.R., Mestre, N.C.C., Pedro, P.Z., Carteny, C.C., Cormier, B., Keiter, S., Jo, M., 2022. Influence of Particle Size on Ecotoxicity of Low-Density Polyethylene Microplastics , with and without Adsorbed Benzo-a-Pyrene , in Clam *Scrobicularia plana*.
- Saha, M., Togo, A., Mizukawa, K., Murakami, M., Takada, H., Zakaria, M.P., Chiem, N.H., Tuyen, B.C., Prudente, M., Boonyatumanond, R., Sarkar, S.K., Bhattacharya, B., Mishra, P., Tana, T.S., 2009. Sources of sedimentary PAHs in tropical Asian waters: Differentiation between pyrogenic and petrogenic sources by alkyl homolog abundance. *Marine Pollution Bulletin* 58, 189–200. <https://doi.org/10.1016/j.marpolbul.2008.04.049>
- Santos, C., Bueno, C., 2020. Biotransformation in the fi sh *Prochilodus lineatus* : An organ-speci fi c approach to cypla gene expression and biochemical activity 80. <https://doi.org/10.1016/j.etap.2020.103467>
- Scott, M.J., Jones, M.N., 2000. The biodegradation of surfactants in the environment. *Biochimica et Biophysica Acta - Biomembranes* 1508, 235–251. [https://doi.org/10.1016/S0304-4157\(00\)00013-7](https://doi.org/10.1016/S0304-4157(00)00013-7)
- Sen, S., Field, J.M., 2013. Genotoxicity of Polycyclic Aromatic Hydrocarbon Metabolites: Radical Cations and Ketones, 1st ed, *Advances in Molecular Toxicology*. Elsevier B.V.

<https://doi.org/10.1016/B978-0-444-62645-5.00003-1>

- Sforzini, A.S., Moore, M.N., Volta, A., Jha, A., Banni, M., 2017. Role of mTOR in autophagic and lysosomal reactions to environmental stressors in molluscs. *Aquatic Toxicology*. <https://doi.org/10.1016/j.aquatox.2017.12.014>
- Trivedi, S.P., Ratn, A., Awasthi, Y., Kumar, M., Trivedi, A., 2021. In vivo assessment of dichlorvos induced histological and biochemical impairments coupled with expression of p53 responsive apoptotic genes in the liver and kidney of fish, *Channa punctatus* (Bloch, 1793). *Comparative Biochemistry and Physiology Part - C: Toxicology and Pharmacology* 245, 109032. <https://doi.org/10.1016/j.cbpc.2021.109032>
- Trivedi, S.P., Singh, S., Trivedi, A., Kumar, M., 2022. Mercuric chloride-induced oxidative stress, genotoxicity, haematological changes and histopathological alterations in fish *Channa punctatus* (Bloch, 1793). *Journal of Fish Biology* 100, 868–883. <https://doi.org/10.1111/jfb.15019>
- Vanzella, T.P., Martinez, C.B.R., Cólus, I.M.S., 2007. Genotoxic and mutagenic effects of diesel oil water soluble fraction on a neotropical fish species. *Mutation Research - Genetic Toxicology and Environmental Mutagenesis* 631, 36–43. <https://doi.org/10.1016/j.mrgentox.2007.04.004>
- WHO, 2003. Polynuclear aromatic hydrocarbons in Drinking-water Background document for development of WHO Guidelines for Drinking-water Quality. *Who/Sde/Wsh/03.04/59* 2, 27.
- Woo, S.J., 2022. Effects of benzo[a]pyrene exposure on black rockfish (*Sebastes schlegelii*): EROD activity, CYP1A protein, and immunohistochemical and histopathological alterations. *Environmental Science and Pollution Research* 29, 4033–4043. <https://doi.org/10.1007/s11356-021-15949-1>
- Yuan, X., Zhao, H., Wang, Y., Wang, L., Li, D., Zhang, A., Yang, X., Ma, X., Yang, D., Zhou, Y., 2021. Expression profile of a novel glutathione S-transferase gene in the marine polychaete *Perinereis aibuhitensis* in short-term responses to phenanthrene, fluoranthene, and benzo [  $\alpha$  ] pyrene. *Marine Pollution Bulletin* 169, 112552. <https://doi.org/10.1016/j.marpolbul.2021.112552>